arXiv:1404.7034v1 [math.OA] 28 Apr 2014

Minimal dynamical systems on connected odd dimensional spaces

Huaxin Lin

Abstract

Let $\beta : S^{2n+1} \to S^{2n+1}$ be a minimal homeomorphism ($n \geq 1$). We show that the crossed product $C(S^{2n+1}) \rtimes_\beta \mathbb{Z}$ has rational tracial rank at most one. More generally, let Ω be a connected compact metric space with finite covering dimension and with $H^1(\Omega, \mathbb{Z}) = \{0\}$. Suppose that $K_i(C(\Omega)) = \mathbb{Z} \oplus G_i$ for some finite abelian group G_i, $i = 0, 1$. Let $\beta : \Omega \to \Omega$ be a minimal homeomorphism. We also show that $A = C(\Omega) \rtimes_\beta \mathbb{Z}$ has rational tracial rank at most one and is classifiable. In particular, this applies to the minimal dynamical systems on odd dimensional real projective spaces. This was done by studying the minimal homeomorphisms on $X \times \Omega$, where X is the Cantor set.

1 Introduction

Let Ω be a compact metric space and let $\alpha : \Omega \to \Omega$ be a minimal homeomorphism. We study the resulting crossed product C^*-algebra $C(\Omega) \rtimes_\alpha \mathbb{Z}$. There are interesting interplays between minimal dynamical systems and the study of C^*-algebras. A classical result of Giordano, Putnam and Skau [7] showed that two Cantor minimal systems are strong orbit equivalent if and only if the associated crossed product C^*-algebras are isomorphic. The C^*-algebra theoretic aspect of their result is indebted to the fact that the crossed product C^*-algebras are unital simple A$\mathbb{T}$-algebras of real rank zero and belong to the class of classifiable C^*-algebras (see Definition 2.8 below). Therefore these C^*-algebras are classified up to isomorphisms by their Elliott invariant, namely, in this case, by their ordered K-theory. In turn, the Cantor minimal systems are classified up to strong orbit equivalence by their induced K-theory. C^*-algebras of the form $C(\Omega) \rtimes_\alpha \mathbb{Z}$ are always simple when α are minimal. These C^*-algebras provide a rich source of unital separable amenable simple C^*-algebras which satisfy the so-called Universal Coefficient Theorem. On the other hand, the rapidly developing Elliott program, otherwise known as the program of classification of amenable C^*-algebras by K-theoretical invariants, provides a possible way to characterize minimal dynamical systems by their K-theoretical invariants. It is therefore important to know when $C(\Omega) \rtimes_\alpha \mathbb{Z}$ belongs to the classifiable class of amenable simple C^*-algebras (in the sense of the Elliott program). Elliott and Evans ([4]) showed that all irrational rotation algebras, which are crossed product C^*-algebras from minimal dynamical systems on the circle, are classifiable. In fact, they are unital A$\mathbb{T}$-algebras of real rank zero. Let $A = C(\Omega) \rtimes_\alpha \mathbb{Z}$. In [19], it is shown that, if Ω has finite covering dimension and $\rho_A(K_0(A))$, the tracial image of $K_0(A)$, is dense in $\mathrm{Aff}(T(A))$, the space of real continuous affine functions on the tracial state space, then A has tracial rank zero (the converse also holds). Consequently A is classifiable. In the case that Ω is connected and (Ω, α) is unique ergodic, this result states that if (Ω, α) has an irrational rotation number, then A is classifiable. In particular, this recovers the previously mentioned case of irrational rotations on the circle. With recent developments in the Elliott program ([16], [29] and [11]), the classifiable C^*-algebras now include the class of C^*-algebras which are rationally of tracial rank at most one. The result of [19] was pushed further by Toms and Winter to great generality: if projections in A separate the tracial states, then A has rational tracial rank zero. Moreover these crossed products are also classifiable by the Elliott invariant (see [27]).

On the other hand, during these developments, minimal dynamical systems on $X \times \mathbb{T}$, where X is the Cantor set and $\mathbb{T}$ is the circle, have been studied (see [14] and [15]). More general cases were also studied in [25]. In both [19] and [27], the Putnam algebra A_x was used as the main bridge. One may view A_x as a large C^*-subalgebra of $A = C(\Omega) \rtimes_\alpha \mathbb{Z}$ by "taking away" one point. In [14] and [15], a smaller C^*-subalgebra is used. That C^*-subalgebra may be viewed as a C^*-subalgebra by "taking away" a circle (an idea of Hiroki Matui). This method seemed to be too special to be useful in more general cases. However, it has been recently adopted by K. Strung [24] to obtain very interesting result about crossed products of certain minimal systems on the odd spheres S^{2n+1}. Strung showed that, by studying the minimal systems of product type $X \times S^{2n+1}$, one can provide examples of non-unique ergodic minimal dynamical systems on the odd spheres whose crossed product C^*-algebras are classifiable. Let A be the crossed product obtained from the minimal system on the odd sphere. It should be noted A may not have rational tracial rank zero. In particular, its projections may not separate the tracial states. Nevertheless, A has rational tracial rank at most one, i.e., $A \otimes U$ has tracial rank at most one for any UHF-algebra U of infinite type. Therefore a more general classification result in [10] can be applied.

In [24], the minimal homeomorphisms on the odd spheres are assumed to be limits of periodic homeomorphisms constructed by "fast approximation-conjugation". In this note we will study the general minimal dynamical systems on the odd spheres as well as on the odd dimensional real projective spaces. We show that crossed product C^*-algebras from any minimal dynamical systems on odd spheres or odd dimensional real projective spaces have rational tracial rank at most one and are classifiable. It should be mentioned that there are no minimal homeomorphisms on even spheres or on the even dimensional real projective spaces because of the existence of fixed points. We actually prove much more general results (see 6.1 and 6.2 below). The methods we used here to study the minimal dynamical systems on the product spaces of the form $X \times \Omega$, where X is the Cantor set and Ω is a connected space are those developed in [14] and [15]. We also use a recent uniqueness theorem (see 4.2 below) from [12]. We continue to use the argument of Strung as well as an embedding result of Winter ([30]). The classification result in [10] is also applied.

This note is organized as follows. Section 2 serves as preliminaries of this note. In Section 3, we study the general minimal dynamical systems on the product spaces $X \times \Omega$, where X is the Cantor set and Ω is a connected compact space. Examples of minimal dynamical systems studied in Section 3 are presented in Section 4. Applications are presented in Section 5.

Acknowledgements This work was done during the author's stay at the Research Center for Operator Algebras at East China Normal University.

2 Preliminaries

Definition 2.1. Let A be a unital C^*-algebra. Denote by $U(A)$ the unitary group of A and $U_0(A)$ the connected component of $U(A)$ containing the identity. Denote by $CU(A)$ the closure of the commutator subgroup of $U_0(A)$. Denote by $T(A)$ the tracial state space of A. We also use $T(A)$ for traces of the form $\tau \otimes \mathrm{Tr}$ on $M_n(A)$ for all integer n, where Tr is the standard (un-normalized) trace on M_n. Denote by $\rho_A : K_0(A) \to \mathrm{Aff}(T(A))$ defined by $\rho_A([p]) = \tau(p)$ for all projections in $M_n(A)$, $n = 1, 2, ...$

Definition 2.2. Let $A = M_n$ and $a \in A$. We use $\det(a)$ for the usual determinant. If $A = C(X) \otimes M_n$, we will often identify A with $C(X, M_n)$. If $f \in A$, we use $\det(f)$ for the function $\det(f)(x)$ in $C(X)$.

Let A be a unital C^*-algebra with $T(A) \neq \emptyset$ and let $u \in U_0(A)$. Let $\{u(t) : t \in [0, 1]\} \subset U_0(A)$

be a piecewise smooth continuous path with $u(0)=u$ and $u(1)=1_A$. Define

$$\tilde{\Delta}_\tau(u(t))=\frac{1}{2\pi i}\int_0^1 \tau(\frac{du(t)}{dt}u(t)^*)dt \text{ for all } \tau\in T(A). \tag{e 2.1}$$

If $u\in CU(A)$, then $\tilde{\Delta}_\tau(u(t))\in\overline{\rho_A(K_0(A))}$. This de La Harp-Skandalis determinant (which is independent of the choice of the path) gives a homomorphism

$$\Delta: U_0(A)/CU(A)\to \mathrm{Aff}(T(A))/\overline{\rho_A(K_0(A))} \tag{e 2.2}$$

(see [26]).

If $u\in U(A)$, we will use $\overline{u}$ for its image in $U(A)/CU(A)$.

Definition 2.3. Let Ω be a compact metric space. Denote by $\mathrm{Homeo}(\Omega)$ the set of all homeomorphisms on Ω equipped with the topology of pointwise convergence. Let $\beta\in\mathrm{Homeo}(\Omega)$. Denote by $\tilde{\beta}: C(\Omega)\to C(\Omega)$ the automorphism defined by $\tilde{\beta}(f)=f\circ\beta^{-1}$ for all $f\in C(\Omega)$. If $F\subset\Omega$, denote by χ_F the characteristic function of F. When F is a clopen set, $\chi_F\in C(\Omega)$.

Lemma 2.4. (Lemma 2.1 of [14]) *Let X be the Cantor set and Ω be a connected compact metric space. Let $\beta\in\mathrm{Homeo}(X\times\Omega)$. Then there is $\gamma\in\mathrm{Homeo}(X)$ and a continuous map $\varphi: X\to\mathrm{Homeo}(\Omega)$ such that $\beta(x,\xi)=(\gamma(x),\varphi_x(\xi))$ for all $(x,\xi)\in X\times\Omega$.*

Proof. Let $p_X: X\times\Omega\to X$ and $p_\Omega: X\times\Omega\to\Omega$ be projection maps such that $p_X(x,\xi)=x$ and $p_\Omega(z,\xi)=\xi$ for all $(x,\xi)\in X\times\Omega$. Fix $(x,\xi)\in X\times\Omega$. Then $\{x\}\times\Omega$ is the connected component of $X\times\Omega$ containing (x,ξ). The homeomorphism β maps it into a connected component containing $\beta(x,\xi)=(x_1,\xi_1)$, $x_1=p_X(\beta(x,\xi))$ and $\xi_1=p_\Omega(\beta(x,\xi))$. Therefore as just mentioned, the component is $\{x_1\}\times\Omega$. Define $\gamma(x)=p_X(\beta(x,\xi))$ for $x\in X$. Since $p_X(\beta(x,\xi))=p_X(\beta(x,\xi'))$, it is a well-defined map. Since β is a homeomorphism, it is also ready to see that $\gamma\in\mathrm{Homeo}(X)$. Fix $x\in X$, then map $\varphi_x(\xi)=p_\Omega(\beta(x,\xi))$ is a homeomorphism from Ω onto Ω. It is easy to check that $\varphi: X\to\mathrm{Homeo}(\Omega)$ is continuous. □

2.5. Let X be the Cantor set and Ω be a connected compact metric space. Then

$$K_0(C(X\times\Omega))=C(X,K_0(C(\Omega))) \text{ and } K_1(X\times\Omega)=C(X,K_1(C(\Omega))),$$

where the group $K_i(C(\Omega))$ is viewed as a discrete space, $i=0,1$. Moreover

$$K_0(C(X\times\Omega))_+=\{f\in C(X,K_0(C(\Omega))): f(x)\in K_0(C(\Omega))_+\}=C(X,K_0(C(\Omega)))_+,$$

Let (X,α) be a Cantor minimal system. Denote

$$K^0(X,\alpha,K_0(C(\Omega)))=C(X,K_0(C(\Omega)))/\{f-f\circ\alpha^{-1}: f\in C(X,K_0(C(\Omega)))\} \tag{e 2.3}$$

equipped with the positive cone

$$K^0(X,\alpha,K_0(C(\Omega)))_+=\{[f]: f\in C(X,K_0(C(\Omega)))_+\}. \tag{e 2.4}$$

Denote

$$\ker(\mathrm{id}_X-\alpha^{-1})_{*i} \;=\; \{f\in C(X,K_i(C(\Omega))): f-f\circ\alpha^{-1}=0\}. \tag{e 2.5}$$

Denote

$$K^i(X,\alpha\times\varphi,K_0(C(\Omega)))=C(X,K_i(C(\Omega)))/\{f-f\circ(\alpha\times\varphi)^{-1}: f\in C(X,K_i(\Omega))\} \tag{e 2.6}$$

with $K^0(X,\alpha\times\varphi,K_0(C(\Omega)))$ equipped with the positive cone

$$K^0(X,\alpha\times\varphi,K_0(C(\Omega)))_+=\{[f]: f\in C(X,K_0(C(\Omega)))_+\}. \tag{e 2.7}$$

For $i=0,1$, denote

$$\ker(\mathrm{id}_{X\times\Omega}-(\alpha\times\varphi)^{-1})_{*i} \;=\; \{f\in C(X,K_i(C(\Omega))): f-f\circ(\alpha\times\varphi)^{-1}=0\}. \tag{e 2.8}$$

Proposition 2.6. *Let (X,α) be a Cantor minimal system, Ω be a connected compact metric space and let $\varphi : X \to \mathrm{Homeo}(\Omega)$. Let $A = C(X\times\Omega)_{\alpha\times\varphi}\mathbb{Z}$. Then there are short exact sequences*

$$0 \to K^0(X,\alpha\times\varphi,K_0(\Omega)) \to K_0(A) \to \ker(\mathrm{id}_{X\times\Omega} - (\alpha\times\varphi)^{-1}_{*1}) \to 0 \ \textit{and} \tag{e 2.9}$$
$$0 \to K^1(X,\alpha\times\varphi,K_1(\Omega)) \to K_1(A) \to \ker(\mathrm{id}_{X\times\Omega} - (\alpha\times\varphi)^{-1}_{*0}) \to 0. \tag{e 2.10}$$

Moreover,

$$\rho_A(K^0(X,(\alpha\times\varphi)^{-1},K_0(\Omega))) = \rho_A(K^0(X,\alpha,\mathbb{Z})). \tag{e 2.11}$$

Furthermore, if $(\varphi_x)_{*i} = \mathrm{id}_{K_i(C(\Omega))}$ for all $x\in X$, $i = 0,1$, then

$$0 \to K^0(X,\alpha\times\mathrm{id},K_0(\Omega)) \to K_0(A) \to K_1(C(\Omega)) \to 0 \ \textit{and} \tag{e 2.12}$$
$$0 \to K^1(X,\alpha\times\mathrm{id},K_1(\Omega)) \to K_1(A) \to K_0(C(\Omega)) \to 0. \tag{e 2.13}$$

Proof. It is clear that (e 2.9) and (e 2.10) follow from the Pimsner-Voiculescu six-term exact sequence directly. To see the second statement, we note that Ω is connected and all tracial states on $K_0(C(\Omega))$ agree with the rank. To see the last statement, we note that

$$K_i(C(X\times\Omega)) = C(X,K_i(C(\Omega))),\ \ i = 0,1.$$

In the case $(\varphi_x)_{*i} = \mathrm{id}_{K_i(C(\Omega))}$, $i = 0,1$, (only) constant elements are invariant under $(\alpha\times\varphi)^{-1}_{*i}$. It follows that

$$\ker(\mathrm{id}_{X\times\Omega} - (\alpha\times\varphi)^{-1}_{*i}) = K_{i+1}(C(\Omega)),\ \ i = 0,1.$$

□

Lemma 2.7. *Let Ω be a compact metric space with $U(C(\Omega)) = U_0(C(\Omega))$ and $\beta\in\mathrm{Homeo}(\Omega)$. Then*

$$\rho_B(K_0(B)) = \rho_A(\imath_{*0}(K_0(C(\Omega)))), \tag{e 2.14}$$

where $B = C(\Omega)\rtimes_\beta\mathbb{Z}$ and $\imath : C(\Omega)\to B$ is the natural embedding. Consequently, in 2.6, if $H^1(\Omega,\mathbb{Z}) = \{0\}$, then

$$\rho_A(K_0(A)) = \rho_A(K^0(X,\alpha,\mathbb{Z})).$$

Proof. The first part follows from Chapter VI of [5] (see also 10.10.5 of [1]). For the second part, we note that X has zero dimension so $H^1(X\times\Omega,\mathbb{Z}) = \{0\}$. Therefore, by (e 2.11),

$$\rho_A(K_0(A)) = \rho_A(K^0(X,(\alpha\times\varphi)^{-1},K_0(C(\Omega)))) = \rho_A(K^0(X,\alpha,\mathbb{Z})). \tag{e 2.15}$$

□

Definition 2.8. Denote by $\mathcal{A}$ the class of unital $\mathcal{Z}$-stable separable simple amenable C^*-algebras which satisfy the Universal Coefficient Theorem and which have rational tracial rank at most one, i.e., $A\otimes U$ has tracial rank at most one, where U is any infinite dimensional UHF-algebra (see [10]). This class of C^*-algebras is classifiable in the sense that, if A, $B\in\mathcal{A}$, then $A\cong B$ if and only if they have the isomorphic Elliott invariant ([10]). This class contains all unital simple AH-algebras with no dimension growth as well as the Jiang-Su algebra. A description of the range of the invariant is presented in [17].

3 The C^*-subalgebra A_x

Definition 3.1. Let (X, α) be a Cantor minimal system and Ω be a connected finite dimensional compact metric space. Fix $x \in X$. Denote by A_x the C^*-subalgebra of $A = C(X \times \Omega) \times_{\alpha\times\varphi} \mathbb{Z}$ generated by $C(X \times \Omega)$ and $uC_0(X \setminus \{x\} \times \Omega)$, where u is the unitary in A which implements the action $\alpha \times \varphi$.

Theorem 3.2. *The C^*-algebra A_x is isomorphic to a unital simple AH-algebra with slow dimension growth.*

Proof. Part of the proof is known. As in the proof of the part (5) of Proposition 3.3 of [14], using groupoid C^*-algebras, A_x is simple, since we assume that $\alpha \times \varphi$ is minimal.

We now assume that the dimension of Ω is d. To show A_x is locally AH, we use an argument of Ian Putnam (3.1 of [22]) and proceed as in the proof of Proposition 3.3 of [14]. Let

$$\mathcal{P}_n = \{X(n, v, k) : v \in V_n, k = 1, 2, ..., k(v)\}$$

be a sequence of Kakutani-Rohlin partitions which gives a Bratteli-Vershik model for (X, σ) (see Theorem 4.2 of [8] or Section 2 of [20]). We also assume the roof sets

$$R(\mathcal{P}_n) = \bigcup_{v\in V} X(n, v, h_n(v))$$

shrink to a single point x. Let A_n be the C^*-algebra generated by $C(X\times\Omega)$ and $uC(\mathcal{R}(\mathcal{P}_n)^c\times\Omega)$. Since $R(\mathcal{P}_{n+1}) \subset R(\mathcal{P}_n)$, $A_n \subset A_{n+1}$, $n = 1, 2, ...$ It is easy see that A_x is the norm closure of the union of all A_n' s. By using a similar argument to Lemma 3.1 of [22], it can be shown that A_n is isomorphic to

$$\bigoplus_{v\in\mathcal{V}_n} M_{h_n(v)} \otimes C(X(n, v, h_n(v))) \otimes C(\Omega) \cong \bigoplus_{v\in\mathcal{V}_n} M_{h_n(v)}(C(Y_{n,v})), \tag{e 3.16}$$

where $Y_{n,v}$ is a compact metric space of covering dimension d. In fact, let

$$p_v = \sum_{k=1}^{h_n(v)} \chi_{X(n,v,k)}$$

for each $v \in V$. It is easy to check that $p_v u(1 - \chi_{R(\mathcal{P}_n)}) = u(1 - \chi_{R(\mathcal{P}_n)})p_v$. It follows that p_v is central in A_n. Put $e_{i,j}(n, v) = u^{i-j}\chi_{X(n,v,j)\times\Omega}$. One that, for each n and v, $\{e_{i,j}(n, v)\}_{i,j}$ $(1 \le i, j \le h_n(v))$ form matrix units in A_n with

$$\sum_{i=1}^{h_n(v)} e_{i,i} = \sum_{i=1}^{h(v)} \chi_{X(n,v,i)} = p_{n,v}.$$

Note that

$$\chi_{X(n,v,h_n(v))} A_n \chi_{X(n,v,h_n(v))} = C(X(n, v, h_n(v))) \times \Omega). \tag{e 3.17}$$

One checks that the C^*-subalgebra generated by $\{e_{i,j}(n, v)\}$ and $\chi_{X(n,v,h_n(v))} A_n \chi_{X(n,v,h_n(v))}$ is isomorphic to $M_{h_n(v)}(C(X(n, v, h_n(v)) \times \Omega))$. Therefore (e 3.16) holds. It follows from Theorem 1.1 of [13] that A_x is a unital simple AH-algebra with no dimension growth.

□

Proposition 3.3. *Let (X,α) be a Cantor minimal system, let $x\in X$, let Ω be a connected compact metric space and $\varphi: X\to \mathrm{Homeo}(\Omega)$ be a continuous map. Then, for $i=0,1$,*

$$K_i(A_x)\cong K_i(C(X\times\Omega))/\{f-f\circ(\alpha\times\varphi)^{-1}_{*i}: f(x)=0, f\in C(X,K_i(C(\Omega)))\}. \quad \text{(e 3.18)}$$

Moreover, the imbedding $\imath: A_x\to A$ gives an affine homeomorphism $\imath_\sharp: T(A)\to T(A_x)$ and gives an order isomorphism $\rho_{A_x}(K_0(A_x))=\rho_A(K^0(X,\alpha,\mathbb{Z}))$. Moreover, if $q\in C(X)$ is a projection, then uqu^* and q are equivalent in A_x.

Proof. Let A_n be in the proof of 3.2, i.e., A_n is generated by $C(X\times\Omega)$ and $uC(R(\mathcal{P})^c\times\Omega)$. There is a natural homomorphism from $K_i(C(X\times\Omega))$ to $K_i(A_n)$ (by the embedding of $C(X\times\Omega)$). By (e 3.16), this homomorphism is surjective. For $i=0,1$, the kernel is

$$\{f-f(\alpha\times\varphi)^{-1}_{*1}: f(y)=0 \text{ for all } y\in R(\mathcal{P}_n),\ f\in K_i(C(X\times\Omega))\}.$$

Thus (e 3.18) holds.

We now prove that, for any projection $q\in C(X)$, uqu^* and q are equivalent in A_x. If $q(x)=0$, then $uq\in A_x$. It follows that $qu^*\in A_x$. Therefore uqu^* and q are equivalent in A_x. Suppose that $q(x)\neq 0$. It suffices to show that

$$[q]-[q\circ(\alpha\times\varphi)^{-1}]=0 \text{ in } K_0(A_x). \quad \text{(e 3.19)}$$

Let $f(y)=[q(y)]-[1_{C(X\times\Omega)}(y)]$ for all $y\in X$. Note that $f\in C(X,K_0(C(\Omega)))$ with $f(x)=0$. Note that $1_{C(X\times\Omega)}-1_{C(X\times\Omega)}\circ(\alpha\times\varphi)^{-1}=0$. It follows that

$$[q]-[q\circ(\alpha\times\varphi)^{-1}]=f-f\circ(\alpha\times\varphi)^{-1}+[1_{C(X\times\Omega)}]-[1_{C(X\times\Omega)}\circ(\alpha\times\varphi)^{-1}]. \quad \text{(e 3.20)}$$

$$\text{(e 3.21)}$$

From (e 3.18), this implies that $[q]-[q\circ(\alpha\times\varphi)^{-1}]=0$ in $K_0(A_x)$. This proves that q and uqu^* are equivalent in A_x.

To show $T(A_x)=T(A)$, it suffices to show that every tracial state $\tau\in T(A_x)$ can be extended to tracial state of A. Let U be a clopen neighborhood of x such that $U,\alpha^{-1}(U),\cdots,\alpha^{-n}(U)$ are mutually disjoint. Let $p=\chi_{U\times\Omega}$. We have shown p, upu^*, u^npu^{*n} are mutually equivalent in A_x and mutually orthogonal. The proof that $T(A_x)=T(A)$ can then proceed exactly the same way as that of part (4) of Proposition 3.3 of [14].

To show that $\rho_{A_x}(K_0(A_x))=\rho_A(K^0(X,\alpha,\mathbb{Z}))$, we first note that we have just proved that the map sending $[\chi_O]$ to $[\chi_O]$ (for clopen sets $O\subset X$) is an embedding from $K^0(X,\alpha,\mathbb{Z})$ into $K_0(A_x)$. Since Ω is connected, the subgroup $K^0(X,\alpha,\mathbb{Z})$ injectively maps into $K^0(X,\alpha\times\varphi,K_0(C(\Omega)))\subset K_0(A)$.

□

4 Tracial rank

Definition 4.1. Let T be a compact Choquet simplex. Suppose that Y is a compact metric space and $L: C(Y)_{s.a.}\to \mathrm{Aff}(T)$ is an affine homomorphism. We say L is unital and strictly positive if $L(1_{C(Y)})(\tau)=1$ for all $\tau\in T$, and $L(f)(\tau)>0$ for all $\tau\in T$ if $f\neq 0$ and $f\geq 0$.

Suppose that $L: C(Y)_{s.a.}\to \mathrm{Aff}(T)$ is a strictly positive affine homomorphism. Let $f\neq 0$ and $f\geq 0$. Then, since $T(A)$ is compact,

$$\inf\{L(f)(\tau): \tau\in T(A)\}>0. \quad \text{(e 4.22)}$$

For each open subset $O \subset Y$, let

$$d(O) = \inf_{\tau \in T}\{\sup\{L(f)(\tau) : 0 \le f \le 1, \text{ supp}(f) \subset O\}\}. \tag{e 4.23}$$

Then, for any non-empty open subset $O \subset Y$, $d(O) > 0$. For each $a \in (0,1)$, let $\{x_1, x_2, ..., x_m\} \subset Y$ be an $a/4$-dense subset. Define

$$D(a,i) = d(B(x_i, a/4)), \ \ i = 1, 2, ..., m. \tag{e 4.24}$$

Put

$$\bigtriangledown_0 (a) = \min\{D(a,i) : i = 1, 2, ..., m\}. \tag{e 4.25}$$

For any $x \in Y$, there exists i such that $B(x,a) \supset B(x_i, a/4)$. Thus

$$d(B(x,a)) \ge (3/4) \bigtriangledown_0 (a). \tag{e 4.26}$$

Put $\bigtriangledown(a) = (3/4)\bigtriangledown_0(a)$ for all $a \in (0,1)$. Now let A be a unital separable simple C^*-algebra with $T(A) = T$ and let $\varphi : C(Y) \to A$ be a unital monomorphism. Then $\varphi_\sharp : C(Y)_{s.a.} \to \text{Aff}(T(A))$ defined by

$$\varphi_\sharp(f)(\tau) = \tau \circ \varphi(f) \ \text{ for all } \ f \in C(Y)_{s.a.}$$

is a unital strictly positive affine homomorphism. It is easy to check that, for any $1 > \sigma > 0$, there is a finite subset $\mathcal{H} \subset C(Y)_{s.a.}$ and $\eta > 0$ such that

$$\mu_{\tau\circ\varphi}(B(x,r)) \ge \bigtriangledown(r) \tag{e 4.27}$$

for all open ball with radius $r \ge \sigma$, provided that

$$|\tau \circ \varphi(g) - L(g)(\tau)| < \eta \ \text{ for all } \ g \in \mathcal{H}, \tag{e 4.28}$$

where $\mu_{\tau\circ\varphi}$ is the Borel probability measure induced by the state $\tau \circ \varphi$.

Let B be a unital C^*-algebra with $T(B) \neq \emptyset$. Note that, by [26], there is a splitting short exact sequence:

$$0 \to \text{Aff}(T(B)) \to \cup_{n=1}^\infty U(M_n(B))/CU(M_n(B)) \to K_1(B) \to 0.$$

Fix a splitting map, we denote by $U_c(K_1(B))$ the image of $K_1(B)$ under the splitting map.

We will use the following uniqueness theorem:

Theorem 4.2. (Theorem 5.9 of [12]) *Let Y be a compact metric space and let T be a compact Choquet simplex. Suppose that $L : C(Y)_{s.a.} \to \text{Aff}(T(A))$ is a unital strictly positive affine homomorphism. Let $\epsilon > 0$ and $\mathcal{F} \subset C(Y)$ be a finite subset. There exists a finite subset $\mathcal{H} \subset C(Y)_{s.a.}$, a finite subset $\mathcal{P} \subset \underline{K}(C(Y))$, a finite subset $\mathcal{U} \subset U_c(K_1(C(Y)))$, $\delta > 0$ and $\eta > 0$ satisfying the following: Suppose that $\varphi_1, \varphi_2 : C(Y) \to A$ are two unital monomorphisms for some unital simple C^*-algebra A of tracial rank at most one with $T(A) = T$ such that*

$$[\varphi_1]|_{\mathcal{P}} = [\varphi_2]|_{\mathcal{P}}, \tag{e 4.29}$$

$$|\tau \circ \varphi_i(g) - L(g)(\tau)| < \delta \ \textit{for all } \ g \in \mathcal{H}, \ i = 1, 2, \ \textit{and} \tag{e 4.30}$$

$$\text{dist}(\overline{\varphi_1(v)\varphi_2(v^*)}, \overline{1_A}) < \eta \ \textit{for all } \ v \in \mathcal{U}. \tag{e 4.31}$$

Then there is a unitary $w \in A$ such that

$$\|w^*\varphi_1(f)w - \varphi_2(f)\| < \epsilon \ \textit{for all } \ f \in \mathcal{F}. \tag{e 4.32}$$

Proof. This follows immediately from Theorem 5.9 of [12] and the discussion above in 4.1. □

The following is well-known.

Lemma 4.3. *Let Y be a compact metric space such that $U(C(Y)) = U_0(C(Y))$. Then, for any $z \in K_1(C(Y))$, there is an integer $m \geq 1$ and a unitary $v \in M_m(C(Y))$ such that*

$$[v] = z \text{ and } \det(v)(y) = 1 \text{ for all } y \in Y.$$

Proof. There exists an integer $m \geq 1$ and a unitary $w \in M_m(C(Y))$ such that $[w] = z$. Put

$$w_{00}(y) = \det(w)(y) \text{ for all } y \in Y. \tag{e 4.33}$$

Then $w_{00} \in U(C(Y)) = U_0(C(Y))$. It follows that

$$w_0(y) = \begin{pmatrix} w_{00}^*(y) & 0 & \cdots & 0 \\ 0 & 1 & \cdots & 0 \\ & & \ddots & \\ & & & 1 \end{pmatrix} \in U_0(M_m(C(Y))). \tag{e 4.34}$$

Define $v = w_0 w$. Then $[v] = [w] = z$ and

$$\det(v)(y) = \det(w_0)(y)\det(w)(y) = 1 \text{ for all } y \in Y. \tag{e 4.35}$$

□

Lemma 4.4. *Let (X, α) be a Cantor minimal system, let Ω be a connected finite dimensional compact metric space with $U(C(\Omega)) = U_0(C(\Omega))$ and let $\varphi : X \to \mathrm{Homeo}(\Omega)$ be a continuous map. Put $A = C(X \times \Omega) \rtimes_{\alpha\times\varphi} \mathbb{Z}$. Suppose that there is $x \in X$ and an integer $k \geq 1$ such that*

$$[\Phi_y] = [\mathrm{id}_{C(\Omega)}] \text{ in } KL(C(\Omega), C(\Omega)), \tag{e 4.36}$$

where $\Phi_y : C(\Omega) \to C(\Omega)$ is defined by

$$\Phi_y(f) = f \circ \varphi^{-1}_{\alpha^{-k+1}(y)} \circ \varphi^{-1}_{\alpha^{-k+2}(y)} \circ \cdots \circ \varphi_y^{-1} \text{ for all } f \in C(\Omega) \tag{e 4.37}$$

and for all $y \in \{\alpha^j(x) : j \in \mathbb{Z}\}$ and α^k is minimal. Let $x \in X$. Then, for any $N \in \mathbb{N}$, $\epsilon > 0$, and any finite subset $\mathcal{F} \subset C(X \times \Omega)$, there is an integer $M > N$, a clopen neighborhood O of x and partial isometry $w \in A_x$ which satisfy the following:

(1) $\alpha^{-N}(O), \alpha^{-N+1}(O), ..., O, \alpha(O), ..., \alpha^M(O)$ are mutually disjoint and $\mu(O) < \epsilon/M$ for every α-invariant probability measure μ;

(2) $w^*w = \chi_O$ and $ww^* = \chi_{\alpha^M(O)}$;

(3) $u^{*i}wu^i \in A_x$ for $i = 0, 1, ..., N-1$;

(4) $\|wf - fw\| < \epsilon$ for all $f \in \mathcal{F}$.

Proof. Since $U(C(\Omega))$ is connected and X has zero-dimensional, one has

$$U(C(X \times \Omega)) = U_0(C(X \times \Omega)). \tag{e 4.38}$$

It follows from 2.7 that

$$\rho_A(K_0(A)) = \rho_A(K^0(X, \alpha, \mathbb{Z})). \tag{e 4.39}$$

Therefore, by 3.3, the embedding $\imath : A_x \to A$ gives

$$\rho_{A_x}(K_0(A_x)) = \rho_A(K_0(A)). \tag{e 4.40}$$

Note A_x is a unital simple AH-algebra with no dimension growth, by 3.2. So $TR(A_x) \le 1$. It is generated by $C(X \times \Omega)$ and $uC((X \setminus \{x\}) \times \Omega)$. The $g \to 1_X \otimes g$ gives a unital embedding from $C(\Omega)$ into $C(X \times \Omega)$. Therefore there is a unital embedding $\imath : C(\Omega) \to A_x$. Let $L : C(\Omega) \to \mathrm{Aff}(T(A_x))$ be the unital strictly positive affine homomorphism induced by $\imath$. Note that $L(g) = \tau(1 \otimes g)$ for all $g \in C(\Omega)_{s.a.}$ and for all $\tau \in T(A_x)$.

Without loss of generality, we may assume that

$$\mathcal{F} = \{f \otimes 1_\Omega, 1_X \otimes g : f \in \mathcal{F}_0 \text{ and } g \in \mathcal{F}_1\},$$

where $\mathcal{F}_0 \subset C(X)$ and $\mathcal{F}_1 \subset C(\Omega)$ are finite subsets. There exists a clopen neighborhood B_x of x such that

$$|f(x) - f(y)| < \epsilon/8 \text{ for all } y \in B_x \text{ and } \quad \text{for all } f \in \mathcal{F}_0. \tag{e 4.41}$$

Since α^k is minimal, we can find $n > N$ such that

$$\alpha^{kn}(x) \subset B_x. \tag{e 4.42}$$

Choose a sufficiently small clopen neighborhood O_x of x such that (1) holds and

$$\alpha^{kn}(y) \in B_x \text{ for all } y \in O_x. \tag{e 4.43}$$

Moreover, by choosing small O_x, we may also require that

$$O_x \cup \alpha^{kn}(O_x) \subset B_x. \tag{e 4.44}$$

Let $p_1 = \chi_{O_x}$ and $q_1 = \chi_{\alpha^{kn}(O_x)}$. Put $M = kn$. Then

$$u^M g p_1 u^{*M} = u^M g u^{*M} q_1 \text{ for all } g \in C(\Omega). \tag{e 4.45}$$

Define

$$\Psi_x = \Phi_{\alpha^{(1-k)(n-1)}(x)} \circ \Phi_{\alpha^{(2-k)(n-2)}(x)} \circ \cdots \circ \Phi_x. \tag{e 4.46}$$

It follows that

$$[\Psi_x] = [\mathrm{id}_{C(\Omega)}] \times [\mathrm{id}_{C(\Omega)}] \times \cdots \times [\mathrm{id}_{C(\Omega)}] = [\mathrm{id}_{C(\Omega)}]. \tag{e 4.47}$$

Let $\mathcal{H} \subset C(\Omega)_{s.a.}$ be a finite subset, $\mathcal{P} \subset \underline{K}(C(\Omega))$ be a finite subset, $\mathcal{U} \subset U_c(K_1(C(\Omega)))$ be a finite subset, $\delta > 0$ and $\eta > 0$ be required by 4.2 for $\epsilon/4$ (in place of ϵ) and $\mathcal{F}_1$ (in place of $\mathcal{F}$) associated with L given above. Let $\mathcal{V} = \{v_1, v_2, ..., v_m\} \subset M_K(C(\Omega))$ such that $\mathcal{U} \subset \{\overline{v_i} : 1 \le i \le m\}$ and, by 4.3,

$$\det(v_i)(y) = 1 \text{ for all } y \in \Omega, \ \ i = 1, 2, ..., m. \tag{e 4.48}$$

There is a finite subset $\mathcal{G} \subset C(\Omega)$ and $\delta_1 > 0$ satisfying the following: Suppose that $h_1, h_2 : C(\Omega) \to B$ are two unital homomorphisms (for any unital C^*-algebra B) such that

$$\|h_1(g) - h_2(g)\| < \delta_1 \text{ for all } g \in \mathcal{G}. \tag{e 4.49}$$

Then

$$[h_1]|_{\mathcal{P}} = [h_2]|_{\mathcal{P}}. \tag{e 4.50}$$

Let $\mathcal{G}_1 = \mathcal{G} \cup \mathcal{H}$ and let

$$U = \mathrm{diag}(\overbrace{u, u, ..., u}^{K}). \tag{e 4.51}$$

There is a neighborhood O of x with $O \subset O_x$ such that

$$\|u^M gpu^{*M} - \Psi_x(g)q\| < \min\{\delta, \delta_1, \epsilon/8\} \text{ for all } g \in \mathcal{G}_1 \text{ and} \tag{e 4.52}$$

$$\|U^M v_i P U^{*M} - (\Psi_x \otimes \mathrm{id}_{M_K})(v_i)Q\| < \eta, \ 1 \le i \le m, \tag{e 4.53}$$

where

$$p = \chi_O, \ q = \chi_{\alpha^M(O)}, P = p \otimes \mathrm{id}_{M_K} \text{ and } Q = q \otimes \mathrm{id}_{M_K}.$$

Define $\psi_{1,0}, \psi_{2,0} : C(\Omega) \to C(X \times \Omega)$ by $\psi_{1,0}(f) = f$ (as constant along X) and $\psi_{2,0}(f) = \Psi_x(f)$ for all $f \in C(\Omega)$. It follows from (e 4.47) that

$$[\psi_{2,0}] = [\psi_{1,0}] \ KL(C(\Omega), C(X \times \Omega)). \tag{e 4.54}$$

Define $\psi_1', \psi_2' : C(\Omega) \to q(C(X \times \Omega))q$ by $\psi_1'(g) = \psi_{1,0}(g)|_{\alpha^M(O)}$ and $\psi_2'(g) = \psi_1' \circ \psi_{2,0}(g) = \Psi_x(g) \cdot q$ for all $g \in C(\Omega)$. It follows from (e 4.54) that

$$[\psi_1'] = [\psi_2'] \text{ in } KL(C(\Omega), qC(X \times \Omega)q). \tag{e 4.55}$$

Denote by j the embedding $qC(X \times \Omega)q \to qA_xq$, $\psi_i = j \circ \psi_i'$, $i = 0, 1$. Then

$$[\psi_1] = [\psi_2] \text{ in } KL(C(\Omega), qA_xq). \tag{e 4.56}$$

It follows from (e 4.52) and 3.2 that

$$|t \circ \psi_1(g) - t \circ \psi_2(g)| < \delta \text{ for all } g \in \mathcal{H} \text{ and for all } t \in T(qA_xq). \tag{e 4.57}$$

Note

$$\tau \circ \psi_1(g) = \tau(q \otimes g) \text{ for all } \tau \in T(A) \text{ and for all } g \in C(\Omega)_{s.a.}. \tag{e 4.58}$$

It follows that

$$L(g)(\tau) = \frac{\tau \circ \psi_1(g)}{\tau(q)} \text{ for all } g \in C(\Omega)_{s.a.} \tag{e 4.59}$$

for all $\tau \in T(A)$.

Note also that $\psi_1(v_i)\psi_2(v_i^*) \in M_K(C(X \times \Omega))$ for some integer $m \ge 1$. By the virtue of Theorem 10 of Chapter VI of [5],

$$\Delta(\psi_1(v_i)\psi_2(v_i^*)) = \Delta(\det(\psi_1(v_i)\psi_2(v_i^*))) \tag{e 4.60}$$

$$= \Delta(\det(\psi_1(v_i))\det(\psi_2(v_i^*))) = \Delta(\psi_1(\det(v_i))\psi_2(\det(v_i^*))) \tag{e 4.61}$$

$$= \Delta(1_{qA_xq}) \in \rho_{A_x}(K_0(qA_xq)). \tag{e 4.62}$$

It follows that

$$\mathrm{dist}(\overline{\psi_1(v_i)\psi_2(v_i^*)}, \overline{1_{qA_xq}}) = 0. \tag{e 4.63}$$

It follows from 4.2 that there is a unitary $w_1 \in qA_xq$ such that

$$\|w_1\psi_2(g)w_1^* - \psi_1(g)\| < \epsilon/4 \text{ for all } g \in \mathcal{F}_1. \tag{e 4.64}$$

There is a unitary normalizer $w_2 \in A_x \cap C^*(X,\alpha)$ of $C(X)$ such that $w_2pw_2^* = q$. Note that w_2 has the form

$$w_2 = \sum_{m\in\mathbb{Z}} u^m \chi_{\Gamma^{-1}(m)},$$

where $\Gamma : X \to \mathbb{Z}$ is a continuous map. Define $\psi_3, \psi_4, \psi_5 : C(\Omega) \to pA_xp$ by $\psi_3(g) = gp$ and $\psi_4(g) = w_2^*w_1u^Mgpu^{*M}w_1^*w_2$ and $\psi_5(g) = w_2^*w_1(g\circ\Psi_x)pw_1^*w_2$ for all $g \in C(\Omega)$. As above, we compute that

$$[\psi_5] = [\psi_3] \text{ in } KL(C(\Omega), pA_xp). \tag{e 4.65}$$

By (e 4.52), the choice of $\mathcal{G}_1$ and δ_1,

$$[\psi_5]|_{\mathcal{P}} = [\psi_4]|_{\mathcal{P}}. \tag{e 4.66}$$

It follows from 3.2 that

$$\tau\circ\psi_3(g) = \tau\circ\psi_4(g) \text{ for all } g \in C(\Omega) \text{ and for all } \tau \in T(pA_xp). \tag{e 4.67}$$

It is clear that

$$\psi_3(v_i)\psi_4(v_i^*) \in CU(pAp). \tag{e 4.68}$$

It follows that

$$\tilde{\Delta}_\tau(\overline{\psi_3(v_i)\psi_4(v_i^*)}) \in \overline{\rho_A(K_0(A))}. \tag{e 4.69}$$

Therefore, by (e 4.40),

$$\mathrm{dist}(\overline{\psi_1(v_i)\psi_2(v^*)}, \overline{1_{qA_xq}}) = 0. \tag{e 4.70}$$

By applying 4.2 again, we obtain a unitary $w_3 \in pA_xp$ such that

$$\|w_3gpw_3^* - \psi_4(g)\| < \epsilon/4 \text{ for all } g \in \mathcal{F}_1. \tag{e 4.71}$$

Put $w = w_2w_3$. Then $w \in A_x$ and

$$w^*w = pw_3^*w_2^*w_2w_3p = p = \chi_O \text{ and } ww^* = w_2w_3w_3^*w_2^* = w_2pw_2^* = q = \chi_{\alpha^M(O)}. \tag{e 4.72}$$

So (2) holds. Moreover (see also (e 4.52)),

$$\begin{aligned}
\|wgpw^* - gq\| &\le \|w_2(w_3gpw_3^*)w_2^* - w_2\psi_4(g)w_2^*\| + \|w_2\psi_4(g)w_2^* - gq\| && \text{(e 4.73)}\\
&< \epsilon/4 + \|w_1u^Mgpu^{*M}w_1^* - gq\| && \text{(e 4.74)}\\
&< \epsilon/4 + \epsilon/8 + \|w_1\Psi_x(g)w_1^* - gq\| && \text{(e 4.75)}\\
&< \epsilon/4 + \epsilon/8 + \epsilon/4 = 5\epsilon/8 \text{ for all } g \in \mathcal{F}_1. && \text{(e 4.76)}
\end{aligned}$$

It follows that

$$\|wg - gw\| = \|wgp - gqw\| = \|(wgp - gqw)w^*\| = \|wgpw^* - gq\| < 5\epsilon/8 \tag{e 4.77}$$

for all $g \in \mathcal{F}_1$. Since $O \cup \alpha^M(O) \subset B_x$, by (e 4.41), for all $f \in \mathcal{F}_0$,

$$\|wf - fw\| \leq \|wpf - wpf(x)\| + \|wpf(x) - f(x)qw\| + \|f(x)qw - fw\| \quad \text{(e 4.78)}$$
$$< \epsilon/8 + \|wf(x) - f(x)w\| + \epsilon/8 = \epsilon/4. \quad \text{(e 4.79)}$$

Thus (4) holds. To see (3), we note that

$$pu^i = pu\chi_{\alpha^{-1}(O)}u\chi_{\alpha^{-2}(O)}\cdots u\chi_{\alpha^{-i}(O)} \text{ and} \quad \text{(e 4.80)}$$
$$(u^{*i}q)^* = qu\chi_{\alpha^{M-1}(O)}u\chi_{\alpha^{M-2}(O)}\cdots u\chi_{\alpha^{M-i}(O)} \quad \text{(e 4.81)}$$

for $i = 1, 2, ..., N-1$. Since $x \in O$, (1) implies that pu^i and $u^{*i}q$ are in A_x. From this one concludes that

$$u^{*i}wu^i \in A_x. \quad \text{(e 4.82)}$$

This proves the lemma. □

In 4.4, if $k = 1$, the assumption (e 4.36) means that

$$[\tilde{\varphi}_x] = [\mathrm{id}_{C(\Omega)}] \text{ in } KL(C(\Omega), C(\Omega)). \quad \text{(e 4.83)}$$

We would like to state the following:

Corollary 4.5. *In 4.4, in the case $k = 1$, the lemma holds if the condition that $U(C(\Omega)) = U_0(C(\Omega))$ is replaced by the following: for each $z \in U(C(\Omega))/U_0(C(\Omega))$, there exists $v \in U(C(\Omega))$ with $[v] = z$ and $h \in C(X)_{s.a.}$ such that*

$$\tilde{\varphi}_y(v) = v\exp(ih(y)) \text{ for all } y \in X. \quad \text{(e 4.84)}$$

Proof. For each $z \in U(C(\Omega))/U_0(C(\Omega))$, choose $v \in U(C(\Omega))$ such that $[v] = z$ and (e 4.84) holds. Therefore the rotation map

$$\Delta_t(v(\alpha \times \varphi)(v^*)) = -t(h) + \rho_{C(X\times\Omega)}(K_0(C(X \times C(\Omega)))) \quad \text{(e 4.85)}$$

for all $t \in T(C(X \times \Omega))$. Since α is minimal, $C(X) \rtimes_\alpha \mathbb{Z}$ is a unital AT-algebra of real rank zero. In particular, $\rho_B(K_0(B))$ is dense in $\mathrm{Aff}(T(B))$, where $B = C(X) \rtimes_\alpha \mathbb{Z}$. Note that

$$\rho_B(K_0(B)) = \rho_A(K^0(X, \alpha, \mathbb{Z})) \text{ and } h(y) \in C(X)_{s.a.}. \quad \text{(e 4.86)}$$

It follows that, for each $\tau \in T(A)$,

$$\Delta_\tau(v(\alpha \times \varphi)(v^*)) \in \overline{\rho_A(K^0(X, \alpha, \mathbb{Z}))}. \quad \text{(e 4.87)}$$

It follows from the last part of 2.6 and (e 4.87) above that

$$\overline{\rho_{A_x}(K_0(A_x))} = \overline{\rho_A(K_0(A))}. \quad \text{(e 4.88)}$$

In the proof of 4.4, if $v_j \in U(C(\Omega))$, then

$$\psi_1(v_j)\psi_2(v_j)^* = v_jq\Psi_x(v_j^*)q = v_j\tilde{\varphi}_x(v_j^*) = \exp(-ih_j) \quad \text{(e 4.89)}$$

for some $h_j \in C(X)_{s.a.}$. It follows that

$$D_\tau(\psi_1(v_j)\psi_2(v_j)) = -\tau(h_jq) \in \overline{\rho_{A_x}(K_0(qA_xq))}, \quad \text{(e 4.90)}$$

since $h_jq \in C(X)$. If $v_j \neq U(C(\Omega))$ and $v_j \in M_K(C(\Omega))$ with $K > 1$, let $w_j = \det v_j \in U(C(\Omega))$ and

$$v_j' = \begin{pmatrix} \det v_j^* & 0 & \cdots & 0 \\ 0 & 1 & \cdots & 0 \\ & & \ddots & \\ & & & 1 \end{pmatrix} v_j.$$

Note that $\det(v_j') = 1$. As in the proof of 4.4, one has

$$\Delta(\psi_1(v_j')\psi_2((v_j')^*)) \in \overline{\rho_{A_x}(K_0(qA_xq))}. \tag{e 4.91}$$

Put $w_j' = (v_j'(v_j^*))^*$. Then

$$\begin{aligned} \Delta(\psi_1(v_j)\psi_2(v_j^*)) &= \Delta(\psi_1(w_j'v_j')\psi_2((w_j'v_j')^*)) && \text{(e 4.92)} \\ &= \Delta(\psi_1(v_j')\psi_2((v_j')^*)\psi_2((w_j')^*)\psi_1(w_j')) && \text{(e 4.93)} \\ &= \Delta(\psi_1(v_j')\psi_2((v_j')^*)) + \Delta(\psi_2((w_j')^*)\psi_1(w_j')) && \text{(e 4.94)} \\ &= \Delta(\psi_1(v_j')\psi_2((v_j')^*)) - \Delta(\psi_1(w_j)\psi_2(w_j^*)) \in \overline{\rho_{A_x}(K_0(qA_xq))}. && \text{(e 4.95)} \end{aligned}$$

Thus (e 4.63) also holds in this. The rest of the proof is exactly the same as that of 4.4. □

The following lemma was taken from the proof of Theorem 5.6 of [14].

Lemma 4.6. *Let (X, α) be a Cantor minimal system, Ω be a compact connected finite dimensional metric space and let $\varphi : X \to \mathrm{Homeo}(\Omega)$ be a continuous map. Suppose that there is $x \in X$ such that, for any $N \in \mathbb{N}$, $\delta > 0$, and any finite subset $\mathcal{F} \subset C(X \times \Omega)$, there is an integer $M > N$, a clopen neighborhood O of x and partial isometry $w \in A_x$ which satisfy the following:*

(1) $\alpha^{-N}(O), \alpha^{-N+1}(O), ..., O, \alpha(O), ..., \alpha^M(O)$ are mutually disjoint and $\mu(O) < \delta/M$ for every α-invariant probability measure μ;

(2) $w^*w = \chi_O$ and $ww^* = \chi_{\alpha^M(O)}$;

(3) $u^{*i}wu^i \in A_x$ for $i = 0, 1, ..., N-1$,

(4) $\|wf - fw\| < \epsilon$ for all $f \in \mathcal{F}$.

Then, for any $\epsilon > 0$, any finite subset $\mathcal{F} \subset A$, there exists a projection $e \in A_x$ satisfies the following:

(a) $\|ea - ae\| < \epsilon$ for all $a \in \mathcal{F}$,

(b) $\mathrm{dist}(pap, eA_xe) < \epsilon$ for all $a \in \mathcal{F}$ and

(c) $\tau(1 - e) < \epsilon$ for all $\tau \in T(A)$.

Proof. It suffices to show the following: for any $\epsilon > 0$, any finite subset $\mathcal{F} \subset C(X \times \Omega)$ and any nonzero element $a \in A_+ \setminus \{0\}$, there exists a projection $e \in A_x \subset A$ such that the following hold:

(1') $\|ef - fe\| < \epsilon$ for all $a \in \mathcal{F} \cup \{u\}$,

(2') $\mathrm{dist}(efe, eA_xe) < \epsilon$ for all $f \in \mathcal{F}$ and

(3') $\tau(1 - e) < \epsilon$ for all $\tau \in T(A)$.

Without loss of generality, we may assume that $\mathcal{F}^* = \mathcal{F}$. Choose $N \subset \mathbb{N}$ so that $2\pi/N < \epsilon$. Put

$$\mathcal{G} = \bigcup_{i=0}^{N-1} u^i \mathcal{F} u^{*i}.$$

We obtain an integer $M > N$, a clopen neighborhood O of x and a partial isometry $w \in A_x$ satisfying (1), (2), (3) and (4).

Put $p = \chi_O$ and $q = \chi_{\alpha^M(O)}$. Define

$$P(t) = p\cos t + w\sin t\cos t + w^*\sin t\cos t + q\sin^2 t \;\; t \in [0, \pi/2]. \tag{e 4.96}$$

Then $P(0) = p$ and $P(\pi/2) = q$. Moreover, one checks that $P(t)$ is a continuous path of projections. By (2), (3) and by the choice of $\mathcal{G}$, one has

$$\|u^{i*}P(t)u^i f - f u^{i*}P(t)u^i\| < \epsilon \tag{e 4.97}$$

for all $t \in [0, \pi/2]$, $i = 0, 1, ..., N-1$ and $f \in \mathcal{F}$. Define

$$e = 1 - \left(\sum_{i=0}^{M-N} u^i p u^{i*} + \sum_{i=1}^{N-1} u^{i*}P(i\pi/2N)u^i\right). \tag{e 4.98}$$

Using (1) and (2), one verifies that e is a projection. By the assumption that $u^{i*}wu^i \in A_x$, $e \in A_x$. By (2) and the fact that

$$\{p, upu^*, u^2pu^{2*}, ..., u^{M-N}p(u^{M-N})^*, u^*P(\pi/2N)u, u^{2*}P(2\pi/2N)u^2, ..., (u^*)^{N-1}P((N-1)\pi/2N)u^{N-1}\}$$

is a set of orthogonal projections, it is ready to verify that

$$\|fe - ef\| < \epsilon \text{ for all } f \in \mathcal{F}. \tag{e 4.99}$$

Since

$$\|P(i\pi/2N) - P((i-1)\pi/2N)\| < \pi/N < \epsilon, \;\; i = 1, 2, ..., N, \tag{e 4.100}$$

one can further verify that

$$\|ue - eu\| < \epsilon. \tag{e 4.101}$$

It is clear that $efe \in A_x$ for all $f \in C(X \times \Omega)$. Note that

$$eue = eu(1-p)e. \tag{e 4.102}$$

Therefore $eue \in A_x$. One also has

$$\tau(1-e) < M\tau(p) < \epsilon \tag{e 4.103}$$

for all $\tau \in T(A)$. □

Lemma 4.7. *Let A be a unital simple C^*-algebra and $B \subset A$, where $1_B = 1_A$ and B is a unital simple AH-algebra with no dimension growth such that $T(B) = T(A)$. Suppose A has the following property: For any $\epsilon > 0$ and any subset $\mathcal{F} \subset A$, there is a projection $e \in B$ such that*

(1) $\|ea - ae\| < \epsilon$ *for all* $a \in \mathcal{F}$,

(2) $\mathrm{dist}(eae, eBe) < \epsilon$ *for all* $a \in \mathcal{F}$ *and*

(3) $\tau(1-e) < \epsilon$ *for all* $\tau \in T(A)$.

Then A has tracial rank at most one.

Proof. We first show that, with the assumption, for any given $d \in A_+ \setminus \{0\}$, one can require that

(3) $1 - e \lesssim d$. We may assume that $0 \le d \le 1$. Put

$$\sigma = \inf\{\tau(d) : \tau \in T(A)\}. \tag{e 4.104}$$

Since A is simple and $T(A)$ is compact, $\sigma > 0$. Choose $\epsilon_0 = \min\{\sigma/2, \epsilon/2\}$. By the assumption, there is a projection $e_1 \in B$ such that

$$\|e_1 d e_1 - d_1\| < \epsilon_0/32 \text{ and } \tau(1-e_1) < \epsilon_0/2 \text{ for all } \tau \in T(A) \tag{e 4.105}$$

for some $d_1 \in e_1 B e_1$. Since

$$\tau(d) = \tau(e_1 d e_1) + \tau((1-e_1)d(1-e_1)) \text{ for all } \tau \in T(A), \tag{e 4.106}$$

$$\|e_1 d e_1\| \geq \sigma - \epsilon_0/2 \geq \sigma/2. \tag{e 4.107}$$

It follows that

$$\|d_1\| \geq \sigma_2 - \epsilon_0/32 \geq 15\sigma/32. \tag{e 4.108}$$

Put $\delta = \epsilon_0/32$. Then, by Proposition 2.2 of [23],

$$0 \neq f_\delta(d_1) \lesssim e_1 d e_1 \sim d^{1/2} e_1 d^{1/2} \lesssim d. \tag{e 4.109}$$

Since B has tracial rank at most one, $e_1 B e_1$ has property (SP). In particular, there is a non-zero projection $e_2 \in \overline{f_\delta(d_1) B f_\delta(d_1)}$. Put

$$\epsilon_2 = \min\{\epsilon/2, \inf\{\tau(e_2) : \tau \in T(A)\}\}. \tag{e 4.110}$$

Then, by the assumption, there is a projection $e \in B$ such that
(1) $\|ea - ae\| < \epsilon_2 \leq \epsilon/2$ for all $a \in \mathcal{F}$,
(2) $\text{dist}(eae, eBe) < \epsilon_2 \leq \epsilon/2$ for all $a \in \mathcal{F}$ and
(3') $\tau(1-e) < \epsilon_2 \leq \tau(e_2)$ for all $\tau \in T(A)$.
Since $T(B) = T(A)$ and B has tracial rank at most one,

$$1 - e \lesssim e_2 \lesssim f_\delta(d_1) \lesssim d. \tag{e 4.111}$$

Note that B has tracial rank at most one. Exactly the same argument used in the proof of Lemma 4.4 of [19] shows that A has tracial rank at most one. Another way to reach the conclusion is to apply Lemma 4.3 of [9]. □

Theorem 4.8. *Let (X, α) be a Cantor minimal system, Ω be a compact connected finite dimensional metric space with $U(C(\Omega)) = U_0(C(\Omega))$ and let $\varphi : X \to \text{Homeo}(\Omega)$ be a continuous map. Suppose that there exists $x \in X$ and an integer $k \geq 1$ such that*

$$[\Phi_y] = [\text{id}_{C(\Omega)}] \text{ in } KL(C(\Omega), C(\Omega)), \tag{e 4.112}$$

where

$$\Phi_y(f) = f \circ \varphi^{-1}_{\alpha^{-k}(y)} \circ \varphi^{-1}_{\alpha^{1-k}(x)} \circ \cdots \circ \varphi^{-1}_{\alpha^{-1}(x)} \circ \varphi^{-1}_x \text{ for all } f \in C(\Omega) \tag{e 4.113}$$

for all $y \in \{\alpha^{j-1}(x) : j \in \mathbb{N}\}$ and suppose that α^k is minimal.

If $\alpha \times \varphi$ is minimal, then $A = C(X \times \Omega) \rtimes_{\alpha\times\varphi} \mathbb{Z}$ has tracial rank at most one. Consequently A is isomorphic to a unital simple AH-algebra with no dimension growth.

Proof. This follows from 4.4, 4.6 and 4.7. □

Theorem 4.9. *Let (X,α) be a Cantor minimal system, Ω be a compact connected finite dimensional metric space and let $\varphi: X\to \mathrm{Homeo}(\Omega)$ be a continuous map. Suppose that*

$$[\varphi_y]=[\mathrm{id}_{C(\Omega)}] \text{ in } KL(C(\Omega),C(\Omega)) \tag{e 4.114}$$

for all $y\in X$ and, for each $z\in U(C(\Omega))/U_0(C(\Omega))$, there exists $v\in U(C(\Omega))$ and $h\in C(X)_{s.a.}$ such that

$$\tilde{\varphi}_y(v)=v\exp(ih(y)) \text{ for all } y\in X. \tag{e 4.115}$$

If $\alpha\times\varphi$ is minimal on $X\times\Omega$, then $A=C(X\times\Omega)\rtimes_{\alpha\times\varphi}\mathbb{Z}$ is isomorphic to a unital simple AH-algebra with no dimension growth.

Proof. This follows from 4.5, 4.6 and 4.7. □

5 Examples

5.1. Let (X,α) be a Cantor minimal system and $\varphi: X\to\mathbb{T}^n$ be a continuous map. One may write $\mathbb{T}^n=\mathbb{R}/\mathbb{Z}\times\mathbb{R}/\mathbb{Z}\times\cdots\times\mathbb{R}/\mathbb{Z}$. Given $\eta\in\mathbb{T}^n$, one write $\eta=(t_1,t_2,...,t_n)$, where $t_j\in\mathbb{R}/\mathbb{Z}$. Define $\eta(\xi)=(s_1+t_1,s_2+t_2,...,s_n+t_n)$, where $\xi=(s_1,s_2,...,s_n)\in\mathbb{T}^n$. Define $(\alpha\times\varphi)(x,\xi)=(\alpha(x),\varphi_x(\xi))$ for all $(x,\xi)\in X\times\mathbb{T}^n$. Since X is totally disconnected, we may also write $\varphi_x=(\exp(i\theta_1(x)),\exp(i\theta_2(x)),...,\exp(i\theta_n(x)))$, where $\theta_j\in C(X)_{s.a.}$. Note for each $x\in X$, $[\varphi_x]=[\mathrm{id}_{C(\Omega)}]$. Let z be the standard unitary generator of $C(\mathbb{T})$. Denote by $z_j\in C(\mathbb{T}^n)$ the function which maps $(s_1,s_2,...,s_n)$ to s_j. Then $\tilde{\varphi}_y(z_j)=z_j\exp(i\theta_j(y))$ for all $y\in X$, $j=1,2,...,n$. Therefore if $\alpha\times\varphi$ is minimal, then 4.9 applies. In particular, when $\alpha\times\varphi$ is minimal, $C(X\times\mathbb{T}^n)\rtimes_{\alpha\times\varphi}\mathbb{Z}$ is a unital simple C^*-algebra with tracial rank at most one. In the case that $n=1$, Lemma 4.2 in [14] provides a necessary and sufficient condition for $\alpha\times\varphi$ being minimal (see also [25]).

5.2. Let $\{m_n\}$ be a sequence of integers with $m_n\ge 2$ and $m_n|m_{n+1}$. Let $\lambda_n:\mathbb{Z}/m_{n+1}\to\mathbb{Z}/m_n$ be the quotient map. The the inverse limit $\varprojlim\mathbb{Z}/m_n$ is the Cantor set. The so-called odometer action α is defined by $\alpha(x)=x+1$ for $x\in\varprojlim\mathbb{Z}/m_n$. Such an action is always minimal. Moreover the family $\{\alpha^k: k\in\mathbb{N}\}$ is equicontinuous on the Cantor set (II.9.6.7 of [3]).

Lemma 5.3. *For each integer $k\ge 2$, there exists an odometer action α on the Cantor set such that α^k is minimal.*

Proof. Fix $k\ge 2$. Choose a sequence of integers $\{m_n\}$ such that $(k,m_n)=1$, i.e., k and m_n are relatively prime and $m_n|m_{n+1}$, $n=1,2,....$ Fix $x\in\varprojlim\mathbb{Z}/m_n\mathbb{Z}$. We will show that $\{\alpha^{mk}(x): m\in\mathbb{N}\}$ is dense. Let $y\in\varprojlim\mathbb{Z}/m_n\mathbb{Z}$. Fix $\epsilon>0$. Since $\{\alpha^m: m\in\mathbb{N}\}$ is equicontinuous, there is $\delta>0$ such that, for any pair of $z_1,z_2\in\varprojlim\mathbb{Z}/m_n\mathbb{Z}$,

$$\mathrm{dist}(\alpha^m(z_1),\alpha^m(z_2))<\epsilon/2 \text{ for all } m\in\mathbb{N}, \tag{e 5.116}$$

provided $\mathrm{dist}(z_1,z_2)<\delta$.

There is an integer $j\ge 1$ and $x',y'\in\mathbb{Z}/m_j\mathbb{Z}$ such that $x_0=\{x'_n\},y_0=\{y'_n\}\in\varprojlim\mathbb{Z}/m_n\mathbb{Z}$ and $x'_n=\gamma_{j,n}(x')$ and $y'_n=\gamma_{j,n}(y')$ for all $n<j$, where $\gamma_{j,n}=\gamma_n\circ\gamma_{n+1}\cdots\circ\gamma_j$ and such that

$$\mathrm{dist}(x_0,x)<\delta \text{ and } \mathrm{dist}(y_0,y)<\delta. \tag{e 5.117}$$

We may assume that $\delta<\epsilon/2$. Since $(k,m_j)=1$, there is $m\in\mathbb{N}$ such that $mk\equiv 1(m_j)$ or $mk\equiv -1(m_j)$. Since $-(m_j-1)\equiv 1(m_j)$, in fact, in both case, there is an integer $l_1\ge 1$ such

that $l_1k \equiv 1(m_j)$. We may assume that $y' = x' + m$ in $\mathbb{Z}/m_j\mathbb{Z}$. Then $y' = x' + ml_1k$ in $\mathbb{Z}/m_j\mathbb{Z}$. Then one computes that

$$\alpha^{ml_1k}(x_0) = x_0 + ml_1k = y_0. \tag{e 5.118}$$

It follows that

$$\begin{aligned} \mathrm{dist}(\alpha^{ml_1k}(x), y) &\leq \mathrm{dist}(\alpha^{ml_1k}(x), \alpha^{ml_1k}(x_0)) + \mathrm{dist}(\alpha^{ml_1k}(x_0), y) && \text{(e 5.119)} \\ &< \epsilon/2 + \mathrm{dist}(y', y) < \epsilon. && \text{(e 5.120)} \end{aligned}$$

□

The following is a result of Karen Strung (Proposition 2.1 and Section 5 of [24]) which we quote here for convenience. Note, if Ω is connected, β^m is minimal for any non-zero integer m.

Proposition 5.4. *Let α be an odometer action on the Cantor set and Ω is a compact metric space. Suppose that $\beta : \Omega \to \Omega$ is a minimal homeomorphism such that β^m is minimal for all $m \in \mathbb{N}$. Then $\alpha \times \beta$ is a minimal homeomorphism on $X \times \Omega$.*

Example 5.5. Let $\beta : S^{2n+1} \to S^{2n+1}$ $(n \geq 1)$ be a minimal homeomorphism. It is known such β exists. Fathi and Herman ([6]) showed that there exists a unique ergodic and minimal diffeomorphism on S^{2n+1}. The group $\mathbb{R}/\mathbb{Z}$ can act on S^{2n+1} freely by rotations. By a result of A. Windsor, there are minimal homeomorphisms β on S^{2n+1} such that β can have any number of ergodic measures ([28]). It follows from 5.4 that $\alpha \times \beta$ are minimal homeomorphisms on $X \times S^{2n+1}$, where α is an odometer which has many invariant probability measures.

Corollary 5.6. *Let α be an odometer on the Cantor set X and (S^{2n+1}, β) be a minimal dynamical system with $n \geq 1$. Then $\alpha \times \beta$ is minimal and $A = C(X \times S^{2n+1}) \rtimes_{\alpha\times\beta} \mathbb{Z}$ has tracial rank at most one.*

Proof. It follows from 5.5 that $\alpha \times \beta$ is minimal. Since β is minimal, it does not have fixed point. Therefore β has zero degree. If follows that $[\beta] = [\mathrm{id}]$ in $KK(C(S^{2n+1}), C(S^{2n+1}))$. Moreover, $U(C(S^{2n+1})) = U_0(C(S^{2n+1}))$. Thus Theorem 4.8 applies (with $k = 1$). □

Example 5.7. Consider an $\mathbb{R}/\mathbb{Z}$ action on RP^{2n+1}. We identify RP^{2n+1} as $SO(2n)$. Define $\gamma : \mathbb{R}/\mathbb{Z} \to SO(2n)$ by

$$\gamma(t) = \begin{pmatrix} \cos(\pi t/2) & \sin(\pi t/2) & 0 & & \\ -\sin(\pi t/2) & \cos(\pi t/2) & 0 & & \\ 0 & 0 & 1 & & \\ & & & \ddots & \\ & & & & 1 \end{pmatrix}. \tag{e 5.121}$$

Define an action $\mathbb{R}/\mathbb{Z} \times SO(2n) \to SO(2n)$ by

$$\Gamma(t)(x) = \gamma(t)x \text{ for all } t \in \mathbb{R}/\mathbb{Z} \text{ and } x \in SO(2n). \tag{e 5.122}$$

It is clear Γ is free and a C^∞ -diffeomorphism. For each $1/n^2 > \delta > 0$ and $r \in \mathbb{Q}/\mathbb{Z}$, by [28], there is a minimal diffeomorphism $\beta_r : SO(2n) \to SO(2)$ such that

$$\mathrm{dist}(\beta_r(x), \gamma(r)x) < \delta \text{ for all } x \in SO(2n). \tag{e 5.123}$$

Corollary 5.8. *Let $n \geq 1$ be an integer. There are odometer actions α on the Cantor set X such that for any minimal homeomorphism β on RP^{2n+1}, $A = C(X \times RP^{2n+1}) \rtimes_{\alpha\times\beta} \mathbb{Z}$ is a unital simple C^*-algebra with tracial rank at most one.*

Proof. First it is well known that $H^1(C(RP^{2n+1}),\mathbb{Z}) = \{0\}$. In other words, $U(C(RP^{2n+1})) = U_0(C(RP^{2n+1}))$.

Note that

$$K_0(C(RP^{2n+1})) = \mathbb{Z} \oplus G_0 \text{ and } K_1(C(RP^{2n+1})) = \mathbb{Z}, \tag{e 5.124}$$

where G_0 is a finite group such that $2g = 0$ for all $g \in G_0$. Any automorphism on $K_0(C(RP^{2n+1}))$ induced by an automorphism on $C(RP^{2n+1})$ has the form

$$\begin{pmatrix} \mathrm{id}_{\mathbb{Z}} & 0 \\ \varphi_{2,1} & \varphi_{2,2} \end{pmatrix}, \tag{e 5.125}$$

where $\varphi_{2,1} : \mathbb{Z} \to G_0$ and $\varphi_{2,2} : G_0 \to G_0$ are homomorphisms, since it sends identity of $C(RP^{2n+1})$ to itself and G_0 is finite. Automorphisms of the form in (e 5.125) is a finite subgroup. Suppose that the order of the group is k_1. Then, for any automorphism $\varphi : C(RP^{2n+1}) \to C(RP^{2n+1})$,

$$\varphi_{*0}^{k_1} = \mathrm{id}_{K_0(C(RP^{2n+1}))}. \tag{e 5.126}$$

We note that $H_0(RP^{2n+1},\mathbb{Q}) = \mathbb{Q}$, $H_{2n+1}(RP^{2n+1},\mathbb{Q}) = \mathbb{Q}$ and $H_i(RP^{2n+1}) = \{0\}$ for all other i. Also $H_0(RP^{2n+1},\mathbb{Z}) = \mathbb{Z}$ and $H_{2n+1}(RP^{2n+1},\mathbb{Z}) = \mathbb{Z}$. Let $\tilde{\beta} : C(RP^{2n+1}) \to C(RP^{2n+1})$ be the isomorphism induced by β. Note that $\beta_* = \mathrm{id}$ on $H_0(RP^{2n+1},\mathbb{Z}) = \mathbb{Z}$ and $\tilde{\beta}_*(1) = \pm 1$ on $H_{2n+1}(RP^{2n+1},\mathbb{Z}) = \mathbb{Z}$. Let

$$L_\beta = \sum_{k\ge 0}(-1)^k Tr(\beta_*|(H_k(RP^{2n+1},\mathbb{Q}))) \tag{e 5.127}$$

$$= Tr(\mathrm{id}|(H_0(RP^{2n+1},\mathbb{Q})) + (-1)^{2n+1}Tr(\beta_*|(H_{2n+1}(RP^{2n+1},\mathbb{Q})))) \tag{e 5.128}$$

be the Lefschetz number. If β is minimal, it does not have fixed point. So $L_\beta = 0$. It follows that $\beta_*(1) = 1$ on $H_{2n+1}(RP^{2n+1},\mathbb{Z})$. It follows that, for any minimal homeomorphism β on RP^{2n+1},

$$(\varphi_\beta)_{*1} = \mathrm{id}_{K_1(C(RP^{2n+1}))}. \tag{e 5.129}$$

We compute that

$$K_0(C(RP^{2n+1},\mathbb{Z}/2\mathbb{Z})) = \mathbb{Z}/2\mathbb{Z} \oplus G_0 \text{ and} \tag{e 5.130}$$

$$0 \to \mathbb{Z}/2\mathbb{Z} \to K_1(C(RP^{2n+1},\mathbb{Z}/2\mathbb{Z})) \to G_0 \to 0. \tag{e 5.131}$$

Let k_2 be the order of $\mathrm{Aut}(\mathbb{Z}/2\mathbb{Z} \oplus G_0)$ (which is finite). One also checks, from the above, $K_1(C(RP^{2n+1}),\mathbb{Z}/2\mathbb{Z})$ is a finite abelian group such that $4x = 0$ for all $x \in K_1(C(RP^{2n+1}),\mathbb{Z}/2\mathbb{Z})$. Let k_3 be the order of $\mathrm{Aut}(K_1(C(RP^{2n+1})))$.

Define $k = k_1 \cdot k_2 \cdot k_3$ which depends on n only. Choose a sequence of integers $\{m_j\}$ such that $m_j | m_{j+1}$ for all j and each m_j is prime relative to k. Then, by 5.3, there are odometer actions α on the Cantor set such that α^k is also minimal.

Now let β be a minimal homeomorphism on RP^{2n+1}. Then, by above,

$$[(\tilde{\beta})^k]|_{K_i(C(RP^{2n+1}))} = \mathrm{id},\ \ i = 0,1 \text{ and } [\tilde{\beta})^k]|_{K_i(C(RP^{2n+1}),\mathbb{Z}/2\mathbb{Z})} = \mathrm{id},\ \ i = 0,1. \tag{e 5.132}$$

Note that

$$KL(C(RP^{2n+1}),C(RP^{2n+1})) = Hom_\Lambda \underline{K}(C(RP^{2n+1}),C(RP^{2n+1})) \tag{e 5.133}$$

It follows from 2.11 of [2] that, to check $[\tilde{\beta}^k] = [\mathrm{id}]$, it suffices to show that (e 5.132) holds since $2g = 0$ for all $g \in G_0$. Therefore

$$[\tilde{\beta}^k] = [\mathrm{id}] \text{ in } KL(C(RP^{2n+1}),C(RP^{2n+1})). \tag{e 5.134}$$

By the assumption we also have that α^k is minimal. Hence 4.8 applies to $\alpha \times \beta$.

□

6 Applications

In this section we consider $A = C(\Omega) \rtimes_\beta \mathbb{Z}$, where Ω is a connected compact metric space and β is a minimal homeomorphism on Ω. Specific examples are the cases that $\Omega = S^{2n+1}$ or $\Omega = RP^{2n+1}$, where $n \geq 1$. It should be noted that there are no minimal homeomorphisms on even spheres or even dimensional real projective spaces. Our results can also apply to other connected spaces.

Theorem 6.1. *Let Ω be a connected compact metric space with finite covering dimension such that $U(C(\Omega)) = U_0(C(\Omega))$ and let $\beta : \Omega \to \Omega$ be a minimal homeomorphism. Suppose that $[\tilde{\beta}^k] = [\mathrm{id}]$ in $KL(C(\Omega), C(\Omega))$ for some integer $k \geq 1$, where $\tilde{\beta}(f) = f \circ \beta^{-1}$ for all $f \in C(\Omega)$. Then $A = C(\Omega) \rtimes_\beta \mathbb{Z}$ has rational tracial rank at most one, i.e., $A \otimes U$ has tracial rank at most one for any infinite dimensional UHF-algebra U. In particular, A is in $\mathcal{A}$.*

Proof. First we note that since Ω has finite covering dimension, it follows from [27] that A has finite nuclear dimension. Let α be an odometer action on the Cantor set such that α^k is also minimal. It follows from 5.4 that $\alpha \times \beta$ is a minimal action. Let $B = C(X \times \Omega) \rtimes_{\alpha\times\beta} \mathbb{Z}$.

It follows from 4.8 that B has tracial rank at most one. Consider the embedding $\imath : A \to B$ that sends $C(\Omega) \to C(X \times \Omega)$ and sends implementing unitary to the implementing unitary in a natural way. Any tracial state τ of B is given by an $\alpha \times \beta$-invariant Borel probability measure. Let τ_0 be the unique tracial state on $C(X) \rtimes_\alpha \mathbb{Z}$ which is given by the α-invariant Borel probability measure. Then each $\alpha\times\beta$ -invariant tracial state on $C(X\times\Omega) = C(X)\otimes C(\Omega)$ has the form $\tau_0 \otimes \tau_1$, where τ_1 is a β-invariant tracial state on $C(\Omega)$. It follows that the map $\imath_\sharp : T(B) \to T(A)$ induced by $\imath$ is a homeomorphism. It follows from 2.7 that $\rho_B(K_0(B)) = \rho_B(K^0(X, \alpha, K_0(C(\Omega))))$. Note also since Ω is connected, $\rho_{C(\Omega)}(K_0(C(\Omega))) = \mathbb{Z}$. Therefore if $\tau_0\otimes\tau_1'$ and $\tau_0\otimes\tau_2'$ are two tracial states then they induce the same state on $K^0(X, \alpha, K_0(C(\Omega)))$. It follows that they induce the same state on $K_0(B)$. It follows from Theorem 4.2 of [30] that $A \otimes U$ has tracial rank at most one. It follows from [18] that $A \otimes U$ has tracial rank at most one for all infinite dimensional UHF-algebras U. Since $A = C(\Omega) \rtimes_\beta \mathbb{Z}$, it satisfies the Universal Coefficient Theorem. Furthermore, by [27], A is $\mathcal{Z}$-stable. Therefore $A \in \mathcal{A}$.

□

Theorem 6.2. *Let Ω be a connected compact metric space with finite covering dimension such that $H^1(\Omega, \mathbb{Z}) = \{0\}$ and $K_i(C(\Omega)) = \mathbb{Z}\oplus G_i$, where G_i is a finite group. Suppose that $\beta : \Omega \to \Omega$ is a minimal homeomorphism. Then $A = C(\Omega) \rtimes_\beta \mathbb{Z}$ has rational tracial rank at most one and is in $\mathcal{A}$.*

Proof. This is a corollary of 6.1. We note that $U(C(\Omega)) = U_0(C(\Omega))$. Therefore it suffices to show that $[\tilde{\beta}^k] = [\mathrm{id}]$ in $KL(C(\Omega))$. Similar to the proof of 5.8, it is easy to see that there exists an integer $k_i \geq 1$ such that $((\tilde{\beta})^{k_i})_{*i} = (\tilde{\beta})^{k_i}_{*i} = \mathrm{id}_{K_i(C(\Omega))}$, $i = 0, 1$.

Let r_i be the order of G_i. For each $1 \leq j \leq (r_i)!$, there exists a short exact sequence

$$0 \to \mathbb{Z}/j\mathbb{Z} \oplus G_i/jG_i \to K_i(C(\Omega), \mathbb{Z}/j\mathbb{Z}) \to G_i^{(j)} \to 0, \tag{e 6.135}$$

where $G_i^{(j)} = \{g \in K_i(C(\Omega)) : jg = 0\}$, $i = 0, 1$. Therefore $K_i(C(\Omega), \mathbb{Z}/j\mathbb{Z})$ is a finite group, $i = 0, 1$. Note

$$[\tilde{\beta}]|_{K_i(C(\Omega),\mathbb{Z}/j\mathbb{Z})} \in \mathrm{Aut}(K_i(C(\Omega), \mathbb{Z}/j\mathbb{Z})). \tag{e 6.136}$$

However, $\mathrm{Aut}(K_i(C(\Omega), \mathbb{Z}/j\mathbb{Z}))$ is a finite group. Therefore, for some $m_{i,j} \geq 1$,

$$[\tilde{\beta}^{m_{i,j}}]|_{K_i(C(\Omega),\mathbb{Z}/j\mathbb{Z})} = \mathrm{id}_{K_i(C(\Omega),\mathbb{Z}/j\mathbb{Z})},\ i = 0, 1.$$

Put

$$k = k_1 \cdot k_2 \cdot \prod_{1\le j\le (r_i)!,\, i=0,1} m_{i,j}. \tag{e 6.137}$$

One checks that

$$(\tilde{\beta}^k)_{*i} = \mathrm{id}_{K_i(C(\Omega))} \text{ and } [\tilde{\beta}^k]|_{K_i(C(\Omega),\mathbb{Z}/j\mathbb{Z})} = \mathrm{id}_{K_i(C(\Omega),\mathbb{Z}/j\mathbb{Z})}, \tag{e 6.138}$$

$j = 1, 2, ..., (r_i)!$, $i = 0, 1$. Since r_i is the order of G_i, by 2.11 of [2],

$$[\tilde{\beta}^k] = [\mathrm{id}_{C(\Omega)}]. \tag{e 6.139}$$

□

Corollary 6.3. *Let β be a minimal homeomorphism on S^{2n+1} ($n \ge 1$). Then $A = C(S^{2n+1}) \rtimes_\beta \mathbb{Z}$ has rational tracial rank at most one and is in $\mathcal{A}$.*

Proof. We have noted that $U(C(S^{2n+1})) = U_0(C(S^{2n+1}))$ and any minimal homeomorphism has the property $[\beta] = [\mathrm{id}]$. So 6.1 applies. □

Corollary 6.4. *Let $\beta_1, \beta_2 : S^{2n+1} \to S^{2n+1}$ ($n \ge 1$) be two minimal homeomorphisms and let $A_i = C(S^{2n+1}) \rtimes_{\beta_i} \mathbb{Z}$, $i = 1, 2$. Then $A_1 \cong A_2$ if and only if $T(A_1) \cong T(A_2)$.*

Proof. One computes, by Pimsner-Voiculescu exact sequence ([21]) that $K_i(A_j) = \mathbb{Z} \oplus \mathbb{Z}$ for $i = 0, 1$ and $j = 1, 2$. One also computes that the order of $K_0(A_j)$ is determined by one copy of $\mathbb{Z}$ from the rank of projections of $M_k(C(S^{2n+1}))$ for all k and $K_0(A_1)$ and $K_0(A_2)$ are unital order isomorphic. Furthermore, all traces agree on $K_0(A_1) = K_0(A_2)$. Therefore their Elliott invariant is determined by $T(A_i)$, $i = 1, 2$. Now, by 6.3, the classification theorem in [10] applies. □

Corollary 6.5. *Let β be a minimal homeomorphism on RP^{2n+1} (for $n \ge 1$). Then $A = C(RP^{2n+1}) \rtimes \mathbb{Z}$ has rational tracial rank at most one and is in $\mathcal{A}$.*

Proof. We have noted that $U(C(RP^{2n+1})) = U_0(C(RP^{2n+1}))$ and $K_0(C(RP^{2n+1})) = \mathbb{Z} \oplus G$, where G is a finite group and $K_1(C(RP^{2n+1})) = \mathbb{Z}$. Thus 6.2 applies. □

Corollary 6.6. *Let β_1 and β_2 be two minimal homeomorphisms on RP^{2n+1} (for $n \ge 1$) and let $A_i = C(RP^{2n+1}) \rtimes_{\beta_i} \mathbb{Z}$, $i = 1, 2$. Then $A_1 \cong A_2$ if and only if*

$$K_1(A_1) \cong K_1(A_2),\ (\beta_1)_* = (\beta_2)_* \text{ on } K_0(C(RP^{2n+1})) \text{ and } T(A_1) = T(A_2). \tag{e 6.140}$$

Proof. By 6.5, it suffices to show that A_1 and A_2 have the same Elliott invariant. The assumption shows that

$$K_0(C(RP^{2n+1}))/\{z - z \circ (\beta_1)_* : z \in K_0(C(RP^{2n+1}))\} \tag{e 6.141}$$

$$\cong\ K_0(C(RP^{2n+1}))/\{z - z \circ (\beta_2)_* : z \in K_0(C(RP^{2n+1}))\} \cong \mathbb{Z} \oplus G_0', \tag{e 6.142}$$

where G_0' is a quotient of $\mathrm{Tor}(K_0(C(RP^{2n+1}))$. Moreover they are order isomorphic. By Pimsner-Voiculescu exact sequence, we may write

$$K_0(A_1) = (\mathbb{Z} \oplus G_0') \oplus \mathbb{Z} \cong K_0(A_2). \tag{e 6.143}$$

Since $H^1(RP^{2n+1}, \mathbb{Z}) = \{0\}$, it follows that

$$\rho_{A_1}(K_0(A_1)) = \rho_{A_1}(\mathbb{Z} \oplus G_0') = \mathbb{Z} \text{ and } \rho_{A_2}(K_0(A_2)) = \mathbb{Z}. \tag{e 6.144}$$

It follows that $K_0(A_1)$ and $K_0(A_2)$ are unital order isomorphic. Since all traces of A_i agree on $K_0(A_i)$, $i = 1, 2$. It follows that A_1 and A_2 have isomorphic Elliott invariant. Thus [10] applies. □

References

[1] B. Blackadar, *K-theory for operator algebras*, Mathematical Sciences Research Institute Publications, 5. Springer-Verlag, New York, 1986. viii+338 pp. ISBN: 0-387-96391-X.

[2] M. Dadarlat and T. Loring, *A universal multicoefficient theorem for the Kasparov groups*, Duke Math. J. **84** (1996), 355–377.

[3] J. de Vries, *Elements of topological dynamics*, Mathematics and its Applications, 257. Kluwer Academic Publishers Group, Dordrecht, 1993. xvi+748 pp. ISBN: 0-7923-2287-8.

[4] G. A. Elliott and E. D. Evans, *The structure of the irrational rotation C^*-algebras*, Ann. of Math. **138** (1993), 477–501.

[5] R. Exel, *Rotation numbers for automorphisms of C^*-algebras,* Pacific J. Math. **127** (1987), 31–89.

[6] A. Fathi and M. Herman, *Existence de diffeomorphismes minimaux*, (French) Dynamical systems, Vol. I Warsaw, pp. 37–59. Asterisque, No. 49, Soc. Math. France, Paris, 1977.

[7] T. Giordano, I. Putnam and C. Skau, *Topological orbit equivalence and C^*-crossed products*, J. Reine Angew. Math. **469** (1995), 51–111.

[8] R. Herman, I. Putnam and C. Skau, *Ordered Bratteli diagrams, dimension groups and topological dynamics*, Internat. J. Math. **3** (1992), 827–864.

[9] S. Hu, H. Lin and Y. Xue, *The tracial topological rank of extensions of C^*-algebras*, Math. Scand. **94** (2004), 125–147.

[10] H. Lin, *Asymptotic unitary equivalence and classification of simple amenable C^*-algebras*, Invent. Math. **183** (2011), 385–450.

[11] H. Lin, *Localizing the Elliott conjecture at strongly self-absorbing C^*-algebras, II, an appendix,* J. Reine Angew. Math., to appear (arXiv:0709.1654).

[12] H. Lin, *Homomorphisms from AH-algebras,* preprint (arXiv:1102.4631).

[13] H. Lin, *On local AH algebras*, Memoir. Amer. Math. Soc., to appear (arXiv:1104.0445).

[14] H. Lin and H. Matui, *Minimal dynamical systems on the product of the Cantor set and the circle*, Comm. Math. Phys. **257** (2005), 425–471.

[15] H. Lin and H. Matui, *Minimal dynamical systems on the product of the Cantor set and the circle. II*, Selecta Math. (N.S.) **12** (2006), 199–239.

[16] H. Lin and Z. Niu, *Lifting KK-elements, asymptotic unitary equivalence and classification of simple C^*-algebras*, Adv. Math. **219** (2008), 1729–1769.

[17] H. Lin and Z. Niu, *The range of a class of classifiable separable simple amenable C^*-algebras*, J. Funct. Anal. **260** (2011), 1–29.

[18] H. Lin and W. Sun, *Tensor products of classifiable C^*-algebras,* preprint, (arXiv:1203.3737).

[19] H. Lin and N. C. Phillips, *Crossed products by minimal homeomorphisms*, J. Reine Angew. Math. **641** (2010), 95–122.

[20] H. Matui, *Approximate conjugacy and full groups of Cantor minimal systems*, Publ. Res. Inst. Math. Sci. **41** (2005), 695–722.

[21] M. Pimsner and D. Voiculescu, *Exact sequences for K-groups and Ext-groups of certain cross-product C^*-algebras*, J. Operator Theory **4** (1980), 93–118.

[22] I. Putnam, *The C^*-algebras associated with minimal homeomorphisms of the Cantor set,* Pacific J. Math. **136** (1989), 329–353.

[23] M. Rørdam, *On the structure of simple C^*-algebras tensored with a UHF-algebra. II*, J. Funct. Anal. **107** (1992), 255–269.

[24] K. Strung, *C^*-algebras of minimal dynamical systems of the product of a Cantor set and an odd dimensional sphere*, prepirnt (arXiv:1403.3136).

[25] W. Sun, *Crossed product C^*-algebras of minimal dynamical systems on the product of the Cantor set and the torus*, J. Funct. Anal., **265** (2013), 1105–1169.

[26] K. Thomsen, *Traces, unitary characters and crossed products by Z*, Publ. Res. Inst. Math. Sci. **31** (1995), 1011–1029.

[27] A. Toms and W. Winter, *Minimal dynamics and the classification of C^*-algebras*, Proc. Natl. Acad. Sci. USA **106** (2009), 16942–16943.

[28] A. Windsor, *Minimal but not uniquely ergodic diffeomorphisms. Smooth ergodic theory and its applications* (Seattle, WA, 1999), 809–824, Proc. Sympos. Pure Math., 69, Amer. Math. Soc., Providence, RI, 2001.

[29] W. Winter, *Localizing the Elliott conjecture at strongly self-absorbing C^*-algebras*, J. Reine Angew. Math., to appear (arXiv:0708.0283)

[30] W. Winter, *Classifying crossed product C^*-algebras*, prepirnt (arXiv:1308.5084).